



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

Smith, Graham & Wermuth, Urs D. (2011) tert-Butylaminium 2-carboxy-4,5-dichlorobenzoate. *Acta Crystallographica. Section E: Structure Reports Online*, 67(9), o2461-o2461.

This file was downloaded from: <http://eprints.qut.edu.au/45704/>

© Copyright 2011 please contact the authors


Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

tert-Butylaminium 2-carboxy-4,5-dichlorobenzoate

Graham Smith and Urs D. Wermuth

Acta Cryst. (2011). **E67**, o2461

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The journal publishes over 4000 structures per year. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

tert-Butylaminium 2-carboxy-4,5-dichlorobenzoate

Graham Smith* and Urs D. Wermuth

Faculty of Science and Technology, Queensland University of Technology, GPO Box 2434, Brisbane, Queensland 4001, Australia
Correspondence e-mail: g.smith@qut.edu.au


Received 18 August 2011; accepted 20 August 2011

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.033; wR factor = 0.094; data-to-parameter ratio = 14.5.

In the structure of the title anhydrous salt, $\text{C}_4\text{H}_{12}\text{N}^+\cdot\text{C}_8\text{H}_3\text{Cl}_2\text{O}_4^-$, the 4,5-dichlorophthalate monoanions have the common 'planar' conformation with the carboxyl groups close to coplanar with the benzene ring and with a short intramolecular carboxylic acid $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond. In the crystal, a two-dimensional sheet structure is formed through aminium $\text{N}-\text{H}\cdots\text{O}_{\text{carboxyl}}$ hydrogen-bonding associations.

Related literature

For structures of 1:1 salts of 4,5-dichlorophthalic acid with acyclic aliphatic amines, see: Mattes & Dorau (1986); Bozkurt *et al.* (2006); Smith & Wermuth (2010*a,b,c*).



Experimental

Crystal data

$\text{C}_4\text{H}_{12}\text{N}^+\cdot\text{C}_8\text{H}_3\text{Cl}_2\text{O}_4^-$
 $M_r = 308.15$
Monoclinic, $P2_1/n$
 $a = 6.1778$ (2) Å
 $b = 12.7158$ (4) Å
 $c = 17.7125$ (7) Å
 $\beta = 96.784$ (4)°

$V = 1381.68$ (8) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.48$ mm⁻¹
 $T = 200$ K
 $0.45 \times 0.26 \times 0.18$ mm

Data collection

Oxford Diffraction Gemini-S CCD-detector diffractometer
Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2010)
 $T_{\min} = 0.977$, $T_{\max} = 0.990$
8677 measured reflections
2719 independent reflections
2307 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.094$
 $S = 0.90$
2719 reflections
188 parameters
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.25$ e Å⁻³
 $\Delta\rho_{\min} = -0.21$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|--|----------|-------------|-------------|---------------|
| $\text{N1A}-\text{H11A}\cdots\text{O21}$ | 0.89 (2) | 2.02 (2) | 2.883 (2) | 164 (2) |
| $\text{N1A}-\text{H12A}\cdots\text{O11}^{\text{i}}$ | 0.91 (2) | 1.88 (2) | 2.784 (2) | 174 (2) |
| $\text{N1A}-\text{H13A}\cdots\text{O12}^{\text{ii}}$ | 0.89 (2) | 1.99 (2) | 2.861 (2) | 167 (2) |
| $\text{O21}-\text{H21}\cdots\text{O12}$ | 0.94 (4) | 1.47 (4) | 2.4021 (19) | 173 (4) |

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x - 1, y, z$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) within *WinGX* (Farrugia, 1999); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON*.

The authors acknowledge financial support from the Australian Research Council and the Faculty of Science and Technology and the University Library, Queensland University of Technology.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5620).

References

- Altomare, A., Cascarno, C., Giacovazzo, A., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
Bozkurt, E., Kartal, I., Odabaşoğlu, M. & Büyükgüngör, O. (2006). *Acta Cryst.* **E62**, o4258–o4260.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Mattes, R. & Dorau, A. (1986). *Z. Naturforsch. B*, **41**, 808–814.
Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Smith, G. & Wermuth, U. D. (2010*a*). *Acta Cryst.* **E66**, o133.
Smith, G. & Wermuth, U. D. (2010*b*). *J. Chem. Crystallogr.* **40**, 207–212.
Smith, G. & Wermuth, U. D. (2010*c*). *Acta Cryst.* **C66**, o374–o380.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2011). E67, o2461 [doi:10.1107/S1600536811034131]

***tert*-Butylaminium 2-carboxy-4,5-dichlorobenzoate**

G. Smith and U. D. Wermuth

Comment

4,5-Dichlorophthalic acid (DCPA) commonly forms 1:1 salts with the acyclic aliphatic amine analogues and with these, low-dimensional hydrogen-bonded structures are usually found, featuring the 'planar' hydrogen phthalate anion e.g with isopropylamine (Smith & Wermuth, 2010*a*), diisopropylamine (Smith & Wermuth, 2010*b*), diethylamine, triethylamine and *n*-butylamine (Smith & Wermuth, 2010*c*), the ammonium and tetra(*n*butyl)ammonium salts (Mattes & Dorau, 1986) and the tetramethylammonium salt (Bozkurt *et al.*, 2006). Our 1:1 stoichiometric reaction of DCPA with *t*-butylamine also gave a 1:1 salt $C_4H_{12}N^+ C_8H_3Cl_2O_4^-$, the title compound and the structure is reported here.

In this structure the common 'planar' DCPA anion is found (Fig. 1) and has the previously described (Smith & Wermuth, 2010*c*) short intramolecular carboxylic acid $O-H\cdots O_{\text{carboxy}}$ hydrogen bond (Table 1) (torsion angles C1–C2–C21–O22 and C2–C1–C11–O11: 176.59 (18) and 175.26 (17) Å respectively). Other structural features common to this 'planar' monoanion are a lengthening of the C1–C11 and C2–C21 bond lengths [1.522 (2) and 1.528 (3) Å] and distortion of the external bond angles at C1 and C2 [C1–C2–C21, 129.57 (15)° and C2–C1–C11, 128.84 (15)°].

Intermolecular aminium $N-H\cdots O(\text{carboxyl})$ hydrogen bonds (Table 1) link the DCPA monoanions across *b* as well as down the *a* axis, forming a two-dimensional sheet structure (Fig. 2).

Experimental

The title compound was synthesized by heating together for 10 min under reflux, 1 mmol quantities of 4,5-dichlorophthalic acid and *t*-butylamine in 50 ml of 50% ethanol–water. Partial evaporation of the solvent gave colourless crystalline plates from which a specimen was cleaved for the X-ray analysis..

Refinement

Hydrogen atoms potentially involved in hydrogen-bonding interactions were located by difference methods and their positional and isotropic displacement parameters were refined. Other H atoms were included at calculated positions [C–H (aromatic) = 0.93 Å or C–H (methyl) = 0.97 Å] and treated as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}\text{C}(\text{aromatic})$ or $1.5U_{\text{eq}}\text{C}(\text{methyl})$.

Figures




Fig. 1. Molecular conformation and atom-numbering scheme for the title salt, with the hydrogen bonds shown as a dashed lines. Non-H atoms are shown as 50% probability displacement ellipsoids.





Fig. 2. A perspective view the two-dimensional sheet structure looking down the sheet, showing hydrogen-bonding associations as dashed lines.

tert-Butylaminium 2-carboxy-4,5-dichlorobenzoate

Crystal data



$$M_r = 308.15$$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$$a = 6.1778 (2) \text{ \AA}$$

$$b = 12.7158 (4) \text{ \AA}$$

$$c = 17.7125 (7) \text{ \AA}$$

$$\beta = 96.784 (4)^\circ$$

$$V = 1381.68 (8) \text{ \AA}^3$$

$$Z = 4$$

$$F(000) = 640$$

$$D_x = 1.481 \text{ Mg m}^{-3}$$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4272 reflections

$$\theta = 3.3\text{--}28.8^\circ$$

$$\mu = 0.48 \text{ mm}^{-1}$$

$$T = 200 \text{ K}$$

Plate, colourless

$$0.45 \times 0.26 \times 0.18 \text{ mm}$$

Data collection

Oxford Diffraction Gemini-S CCD-detector diffractometer

Radiation source: Enhance (Mo) X-ray source graphite

Detector resolution: $16.077 \text{ pixels mm}^{-1}$

ω scans

Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2010)

$$T_{\min} = 0.977, T_{\max} = 0.990$$

8677 measured reflections

2719 independent reflections

2307 reflections with $I > 2\sigma(I)$

$$R_{\text{int}} = 0.027$$

$$\theta_{\max} = 26.0^\circ, \theta_{\min} = 3.4^\circ$$

$$h = -7 \rightarrow 7$$

$$k = -15 \rightarrow 15$$

$$l = -21 \rightarrow 21$$

Refinement

Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.033$$

$$wR(F^2) = 0.094$$

$$S = 0.90$$

2719 reflections

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0608P)^2 + 0.4558P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

188 parameters

$$\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$$

0 restraints

$$\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|-------------|--------------|--------------|----------------------------------|
| Cl1 | 0.77566 (7) | −0.08494 (3) | 0.92701 (3) | 0.0313 (1) |
| Cl2 | 0.29591 (7) | −0.09103 (3) | 0.84148 (3) | 0.0370 (2) |
| O11 | 0.8896 (2) | 0.29012 (10) | 1.01297 (8) | 0.0394 (4) |
| O12 | 0.6289 (2) | 0.40012 (9) | 0.97033 (8) | 0.0353 (4) |
| O21 | 0.2615 (2) | 0.39679 (10) | 0.90835 (10) | 0.0452 (5) |
| O22 | 0.0363 (2) | 0.28642 (11) | 0.84456 (9) | 0.0475 (5) |
| C1 | 0.5836 (3) | 0.21454 (12) | 0.93969 (9) | 0.0218 (5) |
| C2 | 0.3705 (3) | 0.21275 (13) | 0.90024 (9) | 0.0230 (5) |
| C3 | 0.2876 (3) | 0.11656 (13) | 0.87218 (10) | 0.0239 (5) |
| C4 | 0.4066 (3) | 0.02459 (12) | 0.87963 (9) | 0.0233 (5) |
| C5 | 0.6169 (3) | 0.02693 (12) | 0.91703 (9) | 0.0221 (5) |
| C6 | 0.7012 (3) | 0.12081 (13) | 0.94635 (9) | 0.0236 (5) |
| C11 | 0.7102 (3) | 0.30722 (13) | 0.97701 (9) | 0.0268 (5) |
| C21 | 0.2089 (3) | 0.30285 (14) | 0.88234 (11) | 0.0311 (6) |
| N1A | −0.0938 (3) | 0.54459 (13) | 0.90362 (9) | 0.0257 (5) |
| C1A | −0.2152 (3) | 0.58740 (13) | 0.83097 (10) | 0.0255 (5) |
| C2A | −0.0614 (3) | 0.66054 (18) | 0.79557 (12) | 0.0450 (7) |
| C3A | −0.2819 (3) | 0.49408 (16) | 0.77986 (11) | 0.0380 (6) |
| C4A | −0.4120 (3) | 0.64663 (15) | 0.85216 (12) | 0.0373 (6) |
| H3 | 0.14630 | 0.11430 | 0.84740 | 0.0290* |
| H6 | 0.84200 | 0.12160 | 0.97160 | 0.0280* |
| H21 | 0.400 (6) | 0.399 (3) | 0.936 (2) | 0.109 (12)* |
| H11A | 0.016 (3) | 0.5023 (18) | 0.8953 (12) | 0.039 (6)* |
| H12A | −0.036 (4) | 0.5995 (18) | 0.9325 (13) | 0.044 (6)* |
| H13A | −0.185 (4) | 0.5080 (18) | 0.9289 (13) | 0.042 (6)* |
| H21A | 0.06120 | 0.62120 | 0.78220 | 0.0670* |
| H22A | −0.01140 | 0.71450 | 0.83140 | 0.0670* |
| H23A | −0.13660 | 0.69220 | 0.75070 | 0.0670* |
| H31A | −0.37940 | 0.44980 | 0.80390 | 0.0570* |
| H32A | −0.15460 | 0.45460 | 0.77120 | 0.0570* |
| H33A | −0.35380 | 0.51880 | 0.73220 | 0.0570* |

supplementary materials

| | | | | |
|------|----------|---------|---------|---------|
| H41A | −0.50720 | 0.59880 | 0.87410 | 0.0560* |
| H42A | −0.48830 | 0.67810 | 0.80740 | 0.0560* |
| H43A | −0.36510 | 0.70060 | 0.88840 | 0.0560* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|--------------|-------------|-------------|
| Cl1 | 0.0293 (2) | 0.0232 (2) | 0.0394 (3) | 0.0066 (2) | −0.0041 (2) | −0.0007 (2) |
| Cl2 | 0.0298 (3) | 0.0252 (2) | 0.0534 (3) | −0.0039 (2) | −0.0061 (2) | −0.0113 (2) |
| O11 | 0.0394 (8) | 0.0314 (7) | 0.0433 (8) | −0.0070 (6) | −0.0116 (6) | −0.0082 (6) |
| O12 | 0.0367 (7) | 0.0208 (6) | 0.0486 (8) | −0.0044 (5) | 0.0059 (6) | −0.0061 (5) |
| O21 | 0.0321 (8) | 0.0228 (7) | 0.0798 (11) | 0.0057 (6) | 0.0025 (8) | −0.0041 (7) |
| O22 | 0.0377 (8) | 0.0409 (8) | 0.0591 (10) | 0.0146 (6) | −0.0148 (7) | −0.0028 (7) |
| C1 | 0.0250 (8) | 0.0205 (8) | 0.0198 (8) | −0.0024 (6) | 0.0028 (7) | −0.0010 (6) |
| C2 | 0.0228 (8) | 0.0233 (8) | 0.0231 (8) | 0.0017 (6) | 0.0031 (7) | 0.0010 (6) |
| C3 | 0.0180 (8) | 0.0274 (8) | 0.0257 (8) | −0.0004 (7) | 0.0004 (7) | −0.0007 (7) |
| C4 | 0.0231 (8) | 0.0216 (8) | 0.0249 (8) | −0.0033 (6) | 0.0012 (7) | −0.0026 (7) |
| C5 | 0.0223 (8) | 0.0213 (8) | 0.0225 (8) | 0.0018 (6) | 0.0018 (6) | 0.0016 (6) |
| C6 | 0.0204 (8) | 0.0265 (8) | 0.0228 (8) | −0.0016 (6) | −0.0020 (7) | −0.0007 (7) |
| C11 | 0.0311 (9) | 0.0247 (9) | 0.0251 (9) | −0.0068 (7) | 0.0049 (8) | −0.0032 (7) |
| C21 | 0.0316 (10) | 0.0258 (9) | 0.0361 (10) | 0.0067 (7) | 0.0053 (8) | 0.0027 (8) |
| N1A | 0.0236 (8) | 0.0251 (8) | 0.0268 (8) | 0.0001 (7) | −0.0037 (7) | −0.0033 (6) |
| C1A | 0.0245 (9) | 0.0262 (9) | 0.0246 (9) | 0.0016 (7) | −0.0021 (7) | −0.0001 (7) |
| C2A | 0.0437 (12) | 0.0542 (13) | 0.0374 (11) | −0.0117 (10) | 0.0066 (9) | 0.0081 (10) |
| C3A | 0.0385 (11) | 0.0404 (11) | 0.0315 (10) | 0.0050 (9) | −0.0108 (8) | −0.0097 (8) |
| C4A | 0.0328 (10) | 0.0329 (10) | 0.0451 (11) | 0.0079 (8) | 0.0004 (9) | 0.0001 (9) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|------------------------|-------------|------------------------|-------------|
| Cl1—C5 | 1.7251 (17) | C4—C5 | 1.387 (3) |
| Cl2—C4 | 1.7253 (16) | C5—C6 | 1.379 (2) |
| O11—C11 | 1.231 (2) | C3—H3 | 0.9300 |
| O12—C11 | 1.284 (2) | C6—H6 | 0.9300 |
| O21—C21 | 1.307 (2) | C1A—C2A | 1.517 (3) |
| O22—C21 | 1.208 (2) | C1A—C3A | 1.520 (3) |
| O21—H21 | 0.94 (4) | C1A—C4A | 1.515 (3) |
| N1A—C1A | 1.512 (2) | C2A—H21A | 0.9600 |
| N1A—H11A | 0.89 (2) | C2A—H22A | 0.9600 |
| N1A—H13A | 0.89 (2) | C2A—H23A | 0.9600 |
| N1A—H12A | 0.91 (2) | C3A—H31A | 0.9600 |
| C1—C11 | 1.522 (2) | C3A—H32A | 0.9600 |
| C1—C6 | 1.394 (2) | C3A—H33A | 0.9600 |
| C1—C2 | 1.416 (3) | C4A—H41A | 0.9600 |
| C2—C3 | 1.395 (2) | C4A—H42A | 0.9600 |
| C2—C21 | 1.528 (3) | C4A—H43A | 0.9600 |
| C3—C4 | 1.379 (2) | | |
| Cl1...Cl2 | 3.1661 (7) | O11...C21 ^v | 3.217 (2) |
| Cl1...O11 ⁱ | 3.4197 (14) | O11...Cl1 ⁱ | 3.4197 (14) |

| | | | |
|---------------------------|--------------|-------------------------|--------------|
| C11...C3 ⁱⁱ | 3.6461 (18) | O11...N1A ^{vi} | 2.784 (2) |
| Cl2...Cl1 | 3.1661 (7) | O12...C21 | 3.120 (2) |
| C11...H43A ⁱⁱⁱ | 2.9200 | O12...O12 ^{vi} | 3.2387 (17) |
| C11...H6 ⁱ | 2.8300 | O12...O21 | 2.4021 (19) |
| Cl2...H22A ^{iv} | 3.1100 | O12...N1A ^v | 2.861 (2) |
| O11...O22 ^v | 3.219 (2) | O21...C11 | 3.109 (2) |
| C21—O21—H21 | 113 (2) | C4—C3—H3 | 119.00 |
| C1A—N1A—H11A | 112.8 (14) | C1—C6—H6 | 119.00 |
| C1A—N1A—H12A | 108.9 (14) | C5—C6—H6 | 119.00 |
| H11A—N1A—H12A | 107 (2) | C3A—C1A—C4A | 111.52 (15) |
| H11A—N1A—H13A | 108 (2) | N1A—C1A—C2A | 107.49 (15) |
| C1A—N1A—H13A | 109.6 (15) | N1A—C1A—C3A | 107.35 (14) |
| H12A—N1A—H13A | 110 (2) | N1A—C1A—C4A | 107.46 (15) |
| C2—C1—C11 | 128.84 (15) | C2A—C1A—C3A | 111.81 (16) |
| C6—C1—C11 | 112.93 (15) | C2A—C1A—C4A | 110.97 (15) |
| C2—C1—C6 | 118.24 (15) | C1A—C2A—H21A | 109.00 |
| C1—C2—C21 | 129.57 (15) | C1A—C2A—H22A | 109.00 |
| C1—C2—C3 | 118.09 (15) | C1A—C2A—H23A | 109.00 |
| C3—C2—C21 | 112.35 (16) | H21A—C2A—H22A | 109.00 |
| C2—C3—C4 | 122.71 (17) | H21A—C2A—H23A | 109.00 |
| Cl2—C4—C5 | 120.72 (12) | H22A—C2A—H23A | 109.00 |
| Cl2—C4—C3 | 120.19 (14) | C1A—C3A—H31A | 110.00 |
| C3—C4—C5 | 119.08 (15) | C1A—C3A—H32A | 109.00 |
| Cl1—C5—C4 | 121.35 (12) | C1A—C3A—H33A | 109.00 |
| Cl1—C5—C6 | 119.36 (14) | H31A—C3A—H32A | 109.00 |
| C4—C5—C6 | 119.28 (15) | H31A—C3A—H33A | 109.00 |
| C1—C6—C5 | 122.58 (17) | H32A—C3A—H33A | 109.00 |
| O11—C11—C1 | 118.25 (15) | C1A—C4A—H41A | 109.00 |
| O11—C11—O12 | 121.94 (16) | C1A—C4A—H42A | 109.00 |
| O12—C11—C1 | 119.82 (15) | C1A—C4A—H43A | 109.00 |
| O21—C21—C2 | 118.87 (16) | H41A—C4A—H42A | 110.00 |
| O21—C21—O22 | 121.30 (17) | H41A—C4A—H43A | 109.00 |
| O22—C21—C2 | 119.83 (16) | H42A—C4A—H43A | 109.00 |
| C2—C3—H3 | 119.00 | | |
| C6—C1—C2—C3 | 1.9 (2) | C1—C2—C21—O21 | −3.2 (3) |
| C6—C1—C2—C21 | −177.99 (17) | C1—C2—C21—O22 | 176.59 (18) |
| C11—C1—C2—C3 | −178.68 (16) | C3—C2—C21—O21 | 176.90 (17) |
| C11—C1—C2—C21 | 1.4 (3) | C3—C2—C21—O22 | −3.3 (2) |
| C2—C1—C6—C5 | −0.9 (2) | C2—C3—C4—Cl2 | −178.43 (14) |
| C11—C1—C6—C5 | 179.57 (15) | C2—C3—C4—C5 | 0.5 (3) |
| C2—C1—C11—O11 | 175.26 (17) | Cl2—C4—C5—Cl1 | −0.1 (2) |
| C2—C1—C11—O12 | −4.8 (3) | Cl2—C4—C5—C6 | 179.48 (13) |
| C6—C1—C11—O11 | −5.3 (2) | C3—C4—C5—Cl1 | −179.02 (13) |
| C6—C1—C11—O12 | 174.65 (15) | C3—C4—C5—C6 | 0.5 (2) |
| C1—C2—C3—C4 | −1.8 (3) | Cl1—C5—C6—C1 | 179.23 (13) |
| C21—C2—C3—C4 | 178.14 (16) | C4—C5—C6—C1 | −0.3 (3) |

Symmetry codes: (i) $-x+2, -y, -z+2$; (ii) $-x+1, -y, -z+2$; (iii) $x+1, y-1, z$; (iv) $x, y-1, z$; (v) $x+1, y, z$; (vi) $-x+1, -y+1, -z+2$.

Hydrogen-bond geometry (\AA , $^\circ$)

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|--------------------------------------|----------|-------------|-------------|---------------|
| N1A—H11A \cdots O21 | 0.89 (2) | 2.02 (2) | 2.883 (2) | 164 (2) |
| N1A—H12A \cdots O11 ^{vi} | 0.91 (2) | 1.88 (2) | 2.784 (2) | 174 (2) |
| N1A—H13A \cdots O12 ^{vii} | 0.89 (2) | 1.99 (2) | 2.861 (2) | 167 (2) |
| O21—H21 \cdots O12 | 0.94 (4) | 1.47 (4) | 2.4021 (19) | 173 (4) |
| C3—H3 \cdots O22 | 0.93 | 2.29 | 2.671 (2) | 104 |
| C6—H6 \cdots O11 | 0.93 | 2.27 | 2.657 (2) | 104 |

Symmetry codes: (vi) $-x+1, -y+1, -z+2$; (vii) $x-1, y, z$.

Fig. 1




Fig. 2

